arXiv:astro-ph/0505545v1 26 May 2005

Gravitational Wave Astrometry for Rapidly Rotating Neutron Stars and Estimation of Their Distances

Naoki Seto
Theoretical Astrophysics, MC 130-33, California Institute of Technology, Pasadena, CA 91125

We discuss an astrometric timing effect on data analysis of continuous gravitational waves from rapidly rotating isolated neutron stars. Special attention is directed to the possibility of determining their distances by measuring the curvature of the wave fronts. We predict that if continuous gravitational waves from an unknown neutron star with a stable rotation are detected around 1kHz within 1/3yr by initial LIGO detectors and the ellipticity parameter ϵ is smaller than 10^{-6}, the distance r to the source can be estimated with relative error $\Delta r/r$ of $\sim 10\%$ by using the broad band configuration of advanced LIGO detectors over 3 years. By combining the observed amplitude of the waves with the estimated distance, information on the parameter ϵ can be obtained purely through gravitational wave measurements.

PACS numbers: PACS number(s): 95.85.Sz 04.80.Nn, 97.10.Vm

I. INTRODUCTION

Over the last 5 years, several large-scale interferometric gravitational wave detectors, US LIGO [1], Japanese TAMA [2], and British-German GEO [3], were constructed and have given us scientific data. Italian-French VIRGO [4] will be soon in operation. In the next decade, second generation detectors (*e.g.* advanced LIGO [5], LCGT [6]) are expected to be available. More ambitious third generation detectors (*e.g.* EURO [7]) will also be realized in the future. While the current central objective of scientific runs is the direct detection of gravitational waves in 10Hz to a few kHz band, gravitational wave astronomy will be rapidly established with the advent of new and more sensitive detectors [8, 9]. Once gravitational waves from a population of sources are detected with relatively small signal-to-noise ratios (SNRs), follow-on detectors would reveal new astronomical information at high SNRs. This kind of consideration is important for the design and operational strategy in planing next generation detectors.

In this paper, we study an astrometric effect relevant to the data analysis of gravitational waves from rapidly rotating isolated neutron stars and discuss the possibility of estimating their distances. If gravitational waves from a rotating neutron star are detected, the amplitude contains information about the distance r and the ellipticity parameter ϵ through the combination ϵ/r. The parameter ϵ characterizes the non-axisymmetry of the star. However, it is not easy to separate these two quantities individually from gravitational waves. Here, we show that by a long term observation of gravitational waves from a source, we can measure a small phase shift induced by an astrometric effect. This means that a rapidly rotating neutron stars are the ideal targets to apply the long-term astrometric studies with planned ground based detectors. Gravitational waves from merging neutron stars or supernovae would last at most order of seconds [8, 9] and are not suitable for detecting the shift. It is also interesting to understand how the quality of the astronomical information depends on the observational duration. For a very short phenomenon, the only information improved by a long data span is the event rate.

To confirm gravitational waves from a rotating neutron star, we have to integrate the signal for a long time to get a sufficient signal-to-noise ratio [10, 11] (see [12] for a recent observational study with GEO and LIGO). The required computational cost for the data analysis depends strongly on whether the neutron star is known or unknown. If its location is unknown, we need a large number of templates, and the data analysis is a very challenging task [13, 14]. The astrometric effect is larger for a closer object, and we implicitly assume a nearby unknown neutron star as our main target. Once gravitational waves from an unknown neutron star are detected, we might identify an electromagnetic-wave counterpart using information obtained from the gravitational wave measurement, and estimate its distance with various traditional methods (see *e.g.* [15]). But a search for an electromagnetic-wave counterpart would be very difficult for a cold neutron star with a large peculiar velocity and a very weak magnetic field (*e.g.* less than $\sim 10^7$gauss critical for millisecond pulsars [16]).

This paper is organized as follows: In the next section we formulate the phase shift induced by the astrometric effect for a rapidly rotating neutron star, and explain a way to estimate its distance though the phase shift. Then, using the Fisher matrix approach, we obtain an expression for the magnitude of error associated with this estimation. This expression is our central result. In section III we apply it to observational situations with specifications for future projects, such as, advanced LIGO. We examine required conditions for the ellipticity ϵ and the source distance r in order to detect gravitational waves from rotating neutron stars and to determine their distances with the proposed method. Then we statistically study whether there are nearby neutron stars to apply our method. Section IV is devoted to a discussion including a brief summary.

II. FORMULATION

We first study how well we can extract information on the distance from the phase modulation of gravitational waves. Similar arguments have been made for radio pulsar timing analyses [17, 19], but this is the first attempt at applying it to gravitational wave data. Our basic aim is to develop a model relevant for evaluating parameter estimation errors. We do not try to make a detailed model required for actual signal analysis *e.g.* including general relativistic time delays in the solar system that are small and introduce no additional source parameters [11]. In addition, the effects of diurnal rotation are not included here, because we are dealing with the case in which the signal integration is of order a few years. Following Ref.[11], we use the solar system barycenter (SSB) rest frame and investigate the effects induced by the motions of the detector as well as the source. We denote the time dependent position of the detector $\boldsymbol{r}_d$ ($r_d = |\boldsymbol{r}_d| \sim 1\mathrm{AU}$), and assume that the source is moving with a constant velocity $\boldsymbol{v}_{ns}$ as $r\boldsymbol{n}_0 + \boldsymbol{v}_{ns}(t + r/c)$ in the SSB frame. Here $r\boldsymbol{n}_0$ is the position of the source ($\boldsymbol{n}_0$; source direction) from which the emitted wave front reaches the SSB at $t = 0$. We define $\boldsymbol{n}_0 = (\sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$ in the spherical SSB coordinate with θ being the angle between the source direction $\boldsymbol{n}_0$ and the normal vector of the orbital plane (ecliptic) of the Earth. We write the gravitational wave phase Ψ_{SSB} observed at the SSB in the form $\Psi_{SSB}(t) = \Phi_0 + 2\pi\sum_{k=0}^{3} f_k t^{k+1}/(k+1)!$ and perturbatively include the phase modulation due to the detector's and source's motions with two small expansion parameters; $\alpha = |\boldsymbol{r}_d/r|$ and $\beta = |\boldsymbol{v}_{ns}t|/r$. After some algebra, the phase Ψ_d observed by the detector is given to $O(\alpha^m\beta^n)$ $(n + m \leq 2)$ as

$$\begin{aligned}\Psi_d \;=\; & \Phi_0 + 2\pi\sum_{k=0}^{3} f_k \frac{t^{k+1}}{(k+1)!} + \frac{2\pi r}{c}\left(\boldsymbol{n}_0\cdot\frac{\boldsymbol{r}_d}{r} + \frac{\boldsymbol{v}_{ns\perp}t}{r}\cdot\frac{\boldsymbol{r}_d}{r} - \frac{1}{2r^2}(1-(\boldsymbol{n}_0\cdot\boldsymbol{r}_d)^2)\right)\sum_{k=0}^{3} f_k\frac{t^k}{k!} \\ & + \text{higher order terms,} \qquad (2.1)\end{aligned}$$

where $\boldsymbol{v}_{ns\perp}$ is the velocity component of $\boldsymbol{v}_{ns}$ perpendicular to the direction $\boldsymbol{n}_0$ [17]. There are no terms of order $O(\beta)$ and $O(\beta^2)$ [11]. In the large parenthesis in the above equation, the $O(\alpha)$ term is nothing but the plane wave effect that has been analyzed in the literature and can be used to determine the direction of the source. This term is periodic with a $1\mathrm{yr}^{-1}$ frequency. The $O(\alpha\beta)$ term was discussed to some extent for gravitational wave data analysis in Ref.[11, 20] and provides information on the proper motion of the source on the sky. The distance r to the source is estimated through the $O(\alpha^2)$ term. This term probes the curvature of the wave fronts as they deviate from plane wave propagation. In the $O(\alpha^2)$ term, the constant contribution $-1/(2r^2)$ is effectively absorbed in the phase constant Φ_0, while the time dependent oscillating term (with a $2\mathrm{yr}^{-1}$ frequency) is proportional to $\sin^2\theta$. The order of magnitude of the time shift induced by the plane wave $O(\alpha)$ term in one year is written as $\sim r_d\cos\Theta/c \sim 500\mathrm{sec}$ with Θ being the angle between the two vectors $\boldsymbol{n}_0$ and $\boldsymbol{r}_d$. This angle changes by $\Delta\Theta \sim r_d/r$ in one year due to the parallax. Therefore the $O(\alpha^2)$ term $\sim r_d^2/cr \sim 10^{-4}(r/30\mathrm{pc})^{-1}\mathrm{sec}$ can be regarded as a time shift induced by the parallax. Even at $f \sim 1\mathrm{kHz}$, this time shift produces a phase shift much smaller than the 1/4 cycle that is critical for detection of gravitational waves, unless the source is very close to the solar system. Therefore the distance r cannot be estimated without a high SNR observation well beyond the required level for gravitational wave detection.

Using a model $h = h_c\sin[\Psi_d]$ (h_c: constant) for the gravitational wave signal, we calculated the Fisher matrix for the following 10 fitting parameters $\{f_0, f_1, f_2, f_3, \Phi_0, \boldsymbol{v}_{ns,\perp}, r, \theta, \phi\}$ and examined the relative estimation error $\Delta r/r$ with various sets of input parameters as well as the total observational time $T_{obs} \gtrsim 1\mathrm{yr}$. We found that the result $\Delta r/r$ is independent of the parameters ϕ, Φ_0 and inputs $\{f_1, f_2, f_3, f_4, \boldsymbol{v}_{ns\perp}\}$, as long as these inputs are close to reasonable values for real neutron stars. For a integration time T_{obs} longer than $\sim 2\mathrm{yr}$, the resolution $\Delta r/r$ is approximated well by the following expression

$$\frac{\Delta r}{r} = 0.11\left(\frac{r}{100\mathrm{pc}}\right)\left(\frac{SNR}{500}\right)^{-1}\left(\frac{\sin\theta}{\sin\pi/3}\right)^{-2}\left(\frac{f}{1\mathrm{kHz}}\right)^{-1}. \qquad (2.2)$$

We take $\theta = \pi/3$ for a reference value for the angle θ, as it bisects the area of a half sphere $0 \leq \theta \leq \pi/2$. With the specific choice of parameters $\theta = \pi/3$, $f = 1\mathrm{kHz}$, $r = 100\mathrm{pc}$ and $SNR = 500$, the prefactor 0.11 in eq.(2) becomes 9.14, 0.13, 0.109, and 0.107 for $T_{obs} = 1, 2, 3$ and 4 yr respectively. This indicates that for $T_{obs} \gtrsim 2\mathrm{yr}$ the correlation between r and other parameters decreases significantly and becomes nearly stationary. The coefficient becomes 0.107, when we remove f_3 or $\boldsymbol{v}_{ns\perp}$ from the fitting parameters with a fixed observation period $T_{obs} = 3\mathrm{yr}$. In our model we have not directly dealt with the effects of the acceleration of the source. However, these numerical experiments imply that, even if we include acceleration in a relatively simple manner, it would not change the resolution $\Delta r/r$ significantly. When the source is a binary (*e.g.* with its orbital period $\sim 1\mathrm{yr}$), the situation could change considerably.

III. PROSPECTS WITH PROPOSED DETECTORS

Next, we discuss how well we can estimate the distance to a rotating neutron star with the planned gravitational wave detectors, such as, advanced LIGO. We first describe the standard detection criteria of its gravitational wave signal based on SNR for a matched filtering method [10, 13]. Then we use a expression for SNR to evaluate the distance estimation error given by eq.(2.2). For these studies, we pay a close attention to dependence of the ellipticity parameter ϵ and the distance r to the source.

If the rotation axis of the neutron star is identical to one of the principal axes of the inertial moment tensor I_{ij}, the orientation averaged amplitude of gravitational waves becomes $h_c = 8\pi^2 G\sqrt{2/15} I f^2 \epsilon / c^4 r$, where gravitational ellipticity is defined as $\epsilon \equiv (I_{22} - I_{11})/I \ll 1$, and $I = I_{33}$ (x_3: the rotation axis) [8]. With typical numerical values, we obtain

$$h_c = 7.7 \times 10^{-26} \left(\frac{\epsilon}{10^{-8}}\right)\left(\frac{I}{10^{45}\mathrm{g\ cm}^2}\right)\left(\frac{100\mathrm{pc}}{r}\right)\left(\frac{f}{1\mathrm{kHz}}\right)^2. \tag{3.1}$$

Hereafter we fix the inertial moment at $I = 10^{45}\mathrm{g\ cm}^2$ and the gravitational wave frequency at 1kHz. When the rotation axis is different from a principal axis, the star precesses and emits gravitational waves at multiple frequencies [8, 21]. In this paper we use the above equation (3.1) as a reference to relate the characteristic amplitude h_c with two parameters ϵ and r. For the sensitivity of initial and advanced LIGO at 1kHz, we adopt the numerical values given in Figure 1 in Ref.[5] (see also [18]). We take $h_d = 1.5 \times 10^{-22}\mathrm{Hz}^{-1/2}$ for a single 4km detector of initial LIGO and $h_d = 4.2 \times 10^{-24}\mathrm{Hz}^{-1/2}$ for advanced LIGO with a broad band configuration. After averaging with respect to the direction of the source, the effective sensitivity for a monochromatic signal becomes $h_{eff} = h_d\sqrt{2^{-1} 5 T_{obs}^{-1}}$ for an observational period T_{obs}. The factor $2^{-1/2}$ comes from the number of 4km detectors and another factor $5^{1/2}$ is due to the angular average with respect to the source direction. We also study the case for the proposed xylophone-type EURO detector [7] with $h_d = 3.6 \times 10^{-26}\mathrm{Hz}^{-1/2}$ at 1kHz. With these concrete specifications, the signal to noise ratio is calculated as

$$SNR = \frac{h_c}{h_{eff}}, \tag{3.2}$$

and the resolution $\Delta r/r$ is now given as a function of r and ϵ from eqs.(2.2)(3.1) and (3.2).

The characteristic SNR threshold required for detecting an unknown rotating neutron star is discussed in [10] and we take $SNR = 20$ as a reference value for a 1/3yr integration. In figure 1 we plot the $r - \epsilon$ relation for four observational conditions; $SNR = 20$ with (i) initial LIGO and (ii) advanced LIGO for 1/3yr integrations, and $\Delta r/r = 0.1$ with (iii) advanced LIGO and (iv) xylophone-type EURO for 3yr integrations. We have $r \propto \epsilon$ for a given SNR, but $r \propto \epsilon^{1/2}$ for a given resolution $\Delta r/r$. If the parameter ϵ is fixed, we have $SNR \propto r^{-1}$ and $\Delta r/r \propto r^2$.

To detect gravitational waves from neutron stars, a longer integration time is statistically advantageous. This is because of the fact that the amplitude of the detectable signal decreases and the effective survey volume (or equivalently the event rate) increases. However, the number of templates required for the matched filtering method depends strongly on the integration time, and a manageable data length is limited by computational resources. In reality, it might be difficult to deal with data of 1/3 years, especially for a young neutron star with large coefficients f_k ($k \geq 1$) (see eq.(2.1)). In this case, we can detect neutron stars that are closer than the distances shown by the dotted lines in figure 1. However, with a total ~ 10 Teraflops computational power and efficient searching algorithm, the differences between the dotted lines and the maximum detectable distances for initial and advance LIGO would be less than a factor of 2 for most neutron stars (see figure 5 in [13]). Once gravitational wave from a neutron star is confirmed, it would be a relatively easy task to analyze longer data streams with a finer spacing of template parameters. But the available data span would not be much larger than 3 years, considering the time scale of major upgrades to next generation detectors. This is the reason why we studied the case with a 3yr integration for a follow-up analysis of the distance measurement.

In figure 1, the solid line for advanced LIGO and the dotted line for initial LIGO intersect around the interesting value $\epsilon = 10^{-6}$, close to the upper limit from theoretical predictions [10]. Almost all the triangle region in (ϵ, r)-plane bounded by $\epsilon \leq 10^{-6}$ (dashed line) and the dotted line for initial LIGO is below the sold line for advanced LIGO. From viewpoints of future prospects of the gravitational wave astronomy, this fact can be rephrased as follows: if an unknown neutron star is detected around 1kHz with initial LIGO by $\sim 3^{-1}$yr signal integration and the ellipticity has a reasonable magnitude $\epsilon \lesssim 10^{-6}$, advanced LIGO can determine its distance with error of $\sim 10\%$ level within 3 years. This is quite encouraging because blind searches for neutron stars are very hard tasks requiring a huge computational cost, as mentioned earlier [13]. Once it is detected with initial LIGO, advanced LIGO would provide information about its distance, one of the fundamental astronomical parameters for a source. The ellipticity parameter ϵ may



FIG. 1: Relation between the distance r and the gravitational ellipticity ϵ for given observational conditions of an unknown neutron star with gravitational wave frequency $f = 1$kHz. The thin dashed lines represent the maximum distance for detection of gravitational waves with SNR=20 using initial LIGO and advanced LIGO for 1/3yr integrations. The thick solid lines are the maximum distance for resolution $\Delta r/r = 0.1$ through the timing parallax method, shown for advanced LIGO and xylophone-type EURO for 3yr integrations. The dashed line $\epsilon = 10^{-6}$ is an upper limit of theoretical predictions for the ellipticity parameter.

then also be found through knowledge of the amplitude h_c. Both parameters are thus determined purely through gravitational wave measurements. Note that advanced LIGO has the potential to measure the distance to an unknown neutron star that cannot be detected by initial LIGO. The effective sensitivity of the single EURO xylophone-type detector is 82 times better than advanced LIGO detector at 1kHz, and the distance r for a given resolution $\Delta r/r$ is 9.1 times larger.

We have discussed the case in which the gravitational wave data would be integrated without a break (100% duty cycle) for 3 years or so. But our result for $\Delta r/r$ would be approximately valid for a coherent analysis of intermittent data streams that are sampled with appropriate intervals ($\ll 1$yr) and in a sufficient span (*e.g.* 3 years). If the narrow band configuration is available at the target frequency of a source, we can reach the same resolution $\Delta r/r$ with a shorter total integration time. For example, the sensitivity h_d could be improved by a factor ~ 3 at 1kHz with the narrow band configuration of advanced LIGO, and the total integration time could be a factor ~ 10 times smaller than for its broad band configuration.

From Figure 1, we can estimate the distance r for the required resolution $\Delta r/r \sim 0.1$ as a function of the parameter ϵ. Here let us statistically study whether there is at least one rapidly rotating pulsar within this distance. For this purpose we use the argument originally made by Blandford assuming a population of neutron stars whose spin evolutions are primarily determined by energy loss through gravitational radiation [8, 10]. We define τ_B^{-1} as the formation rate of the population in our Galaxy with an initial gravitational wave frequency higher than 1kHz. The magnitude of the time τ_B is highly uncertain but should be much larger than $\tau_B \sim 100$yr corresponding to the Galactic supernova rate. We first evaluate the minimum distance r_{min} to a pulsar of this population, and then compare r_{min} with the distance allowed for the resolution $\Delta r/r = 0.1$. If these two distances coincide, we can expect to have at least one pulsar with resolution $\Delta r/r$ better than 0.1. Our goal here is to calculate the required formation rate τ_B for this condition as a function of the ellipticity ϵ.

In Blandford's model, the amplitude h_c from a neutron star with a fixed distance becomes larger for a larger ϵ. On the other hand, the evolution time scale τ_{GW} becomes smaller for a larger ϵ, and the smallest distance r_{min} would be larger. Here the evolution time scale τ_{GW} is defined as

$$\tau_{GW} = \frac{f}{\dot{f}} = \frac{5c^5}{32\pi^4 GI\epsilon^2 f^4} = 1.8\times 10^8 \left(\frac{\epsilon}{10^{-8}}\right)^{-2}\left(\frac{f}{1\mathrm{kHz}}\right)^{-4} \mathrm{yr}. \tag{3.3}$$

The gravitational frequency becomes smaller by $\sim 30\%$ within a single τ_{GW}. The time scale τ_{GW} becomes the age of the universe $\sim 10^{10}$yr at $\epsilon \sim 10^{-9}(f/1\mathrm{kHz})^{-2}$. For the spatial distribution of the population in our Galaxy, we assume a uniform density in a cylinder with radius $R_{ns} = 10$kpc and height $H_{ns} = 1$kpc. If the parameter ϵ is in the range $10^{-9}(f/1\mathrm{kHz})^2 < \epsilon < 10^{-6}(\tau_B/200\mathrm{yr})^{-1/2}(f/1\mathrm{kHz})^{-2}$, we have

$$r_{min} = \left(3V\tau_B\tau_{GW}^{-1}/4\right)^{1/3} = 44\left(\frac{\tau_B}{200\mathrm{yr}}\right)^{1/3}\left(\frac{\epsilon}{10^{-8}}\right)^{2/3}\left(\frac{f}{1\mathrm{KHz}}\right)^{4/3}\mathrm{pc} \tag{3.4}$$

for the smallest distance. The distance r_{min} becomes larger than the height H_{ns} for $\epsilon > 10^{-6}(\tau_B/200\mathrm{yr})^{-1/2}(f/1\mathrm{kHz})^{-2}$. In this case we have $r_{min} \sim R_{ns}(\tau_B\tau_{GW}^{-1}) \propto \epsilon^{1/2}$. With these expressions and results presented in figure 1, we can now estimate the required formation rate τ_B for observing at least one pulsar with resolution $\Delta r/r = 0.1$. For 3yr advanced LIGO observation at 1kHz, we have $\tau_B < 84(\epsilon/10^{-8})^{-1/2}$yr in the range $10^{-9} < \epsilon < 10^{-5}$. Therefore, a very small τ_B (or a large formation rate τ_B^{-1}) is required from this rough statistical argument. In the case of 3yr EURO-xylophone-type data, we obtain $\tau_B < 6.3\times 10^4(\epsilon/10^{-8})^{-1/2}$yr for $10^{-9} < \epsilon < 10^{-7}$.

So far, we have studied the case with $\Delta r/r \ll 1$. Here, we discuss the opposite case in which the distance to a source is large and we cannot detect the signature of the parallax timing shift. In the case of usual angular parallax measurement, the amplitude of the apparent annual motion of a source on the sky is inversely proportional to its distance r. Therefore, it is straightforward to fit distance with a form r^{-1} in the parameter estimation. We can understand this from the original meaning of the distance unit "parsec" that represents 1/arcsec. The situation is similar to the fitting for the parallax timing. Note also that the error $\Delta(r^{-1}) \sim (\Delta r)/r^2$ for r^{-1} does not depend on the distance r itself, as shown in eq.(2). In the low resolution limit with a fitting result $|(r^{-1})_{fit}| \sim \Delta(r^{-1})$, we can set a constraint for the value r^{-1} in the form $r^{-1} < \max\{(r^{-1})_{fit}, 0\} + N\Delta(r^{-1})$ with $N-\sigma$ statistical significance. At the end we get a lower limit of the distance r as

$$r > (N+x)^{-1}\{\Delta(r^{-1})\}^{-1} = 220\left(\frac{SNR}{500}\right)\left(\frac{\sin\theta}{\sin(\pi/3)}\right)^2\left(\frac{f}{1\mathrm{kHz}}\right)\left(\frac{N+x}{4}\right)^{-1}\mathrm{pc} \tag{3.5}$$

for $T_{obs} \gtrsim 2$yr with $x \equiv \max\{(r^{-1})_{fit}/\Delta(r^{-1}), 0\}$. We can also set a lower limit to the parameter ϵ by using the gravitational wave amplitude h_c.

IV. DISCUSSIONS

In our analysis, the phase $\Phi_{SSB}(t)$ at the SSB is assumed to be simple and be described by a low-order Taylor expansion. Even though many pulsars are known to be extremely accurate clocks, the phase Φ_{SSB} could have a deviation from this simple expression. The origin of these timing deviations can be due to the intrinsic evolution of the pulsars or other effects (*e.g.* the low frequency gravitational wave background). If the deviation is large, especially around the frequency $\sim 2\mathrm{yr}^{-1}$, it would hamper the estimation of the distance through parallax timing even with a high SNR measurement. Here, we briefly describe the actual radio observation of certain pulsars following [22]. PSR B1855+09 is a binary millisecond pulsar with spin period 5.36 msec and orbital period 12.3 day, and PSR B1937+21 is a single millisecond pulsar with spin period 1.56 msec. They are considered to be recycled pulsars. Timing analyses for both pulsars have been performed with data taken at Arecibo Observatory in a ~ 8 year span. A systematic timing deviation of order a few μsec is indicated for PSR B1937+21, but not for PSR B1855+09. The timing residual of the former is expected to be dominated by the intrinsic evolution of the pulsar and not by other effects, such as, the low frequency gravitational wave background. As for the estimated distances through the timing parallax, PSR B1855+09 has a fitted value $\pi \equiv (1\mathrm{kpc}/r)\mathrm{mas} = 1.1 \pm 0.3$mas corresponding to $r = 0.9^{+0.4}_{-0.2}$kpc (1σ), and PSR B1937+21 has a bound $\pi < 0.28$mas (2σ) corresponding to $r > 3.6$kpc. Their distances are also estimated from dispersion measures together with a model of the Galactic distribution of interstellar free electrons. They are 0.7kpc for PSR B1855+09 and 3.6kpc for PSR B1937+21 with uncertainties of $\sim 25\%$, and are consistent with the estimation through the timing parallax. Therefore, the effects of the timing deviation would be moderate in these cases.

Now we summarize this paper. We have discussed an astrometric effect for the analysis of gravitational waves that would allow us to determine the distance to a rapidly rotating isolated neutron star around $f \sim 1$kHz. The resolution

$\Delta r/r$ improves significantly, if the integration period of the signal is longer than ~ 2 years. By using advanced LIGO detectors for 3 years at its designed sensitivity in the broad band configuration, we can estimate its distance with an error of $\Delta r/r \lesssim 0.1$ up to $r \sim 150(\epsilon/10^{-7})^{1/2}$pc, if its spin evolution is relatively simple.

Acknowledgments: The author is grateful to Teviet Creighton for valuable discussions. He also thanks Sterl Phinney, Kip Thorne and Yi Pan for their help and useful conversations, and Asantha Cooray and Steve Furlanetto for carefully reading the manuscript. This work was supported by NASA grant NNG04GK98G and the Japan Society for the Promotion of Science.

[1] A. Abramovici *et al.*, Science **256**, 325 (1992).
[2] M. Ando *et al.* [TAMA Collaboration], Phys. Rev. Lett. **86**, 3950 (2001) [arXiv:astro-ph/0105473].
[3] B. Willke *et al.*, Class. Quant. Grav. **19**, 1377 (2002).
[4] A. Giazotto, Nucl. Instrum. Meth. A **289**, 518 (1990).
[5] E. Gustafson, D. Shoemaker, K. Strain and R. Weiss, 1999, LSC White Paper on Detector Research and Development (LIGO project document T990080-00-D)
[6] K. Kuroda *et al.*, Class. Quant. Grav. **20**, S871 (2003).
[7] EURO homepage: http:/www.astro.cf.ac.uk/geo/euro
[8] K. S. Thorne, in *Three hundred years of gravitation*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, 1987), pp. 330–458.
[9] B. F. Schutz, Class. Quant. Grav. **16**, A131 (1999) [arXiv:gr-qc/9911034].
[10] P. R. Brady, T. Creighton, C. Cutler and B. F. Schutz, Phys. Rev. D **57**, 2101 (1998) [arXiv:gr-qc/9702050].
[11] P. Jaranowski, A. Krolak and B. F. Schutz, Phys. Rev. D **58**, 063001 (1998) [arXiv:gr-qc/9804014].
[12] B. Abbott *et al.* [LIGO Scientific Collaboration], Phys. Rev. D **69**, 082004 (2004) [arXiv:gr-qc/0308050].
[13] P. R. Brady and T. Creighton, Phys. Rev. D **61**, 082001 (2000) [arXiv:gr-qc/9812014].
[14] P. Jaranowski and A. Krolak, Phys. Rev. D **61**, 062001 (2000) [arXiv:gr-qc/9901013]; C. Cutler, I. Gholami and B. Krishnan, arXiv:gr-qc/0505082;
[15] D. A. Frail and J. M. Weisberg, Astron. J. **100**, 743 (1990); G. C. Gomez, R. A. Benjamin, D. P. Cox and P. Donald, Astron. J. **122**, 908 (2001).
[16] E. S. Phinney and S. R. Kulkarni, Annu. Rev. Astron. Astrophys. 32, 591 (1994).
[17] R. Blandford, R. W. Romani and R. Narayan, Journal of Astrophysics and Astronomy, **5**, 369 (1984); T. Damour and N. Deruelle, Ann. Inst. Henri Poincare **44**, 263 (1986); D. C. Becker and R. W. Hellings, Annu. Rev. Astron. Astrophys. 24, 537 (1986).
[18] A. Buonanno and Y. b. Chen, Phys. Rev. D **64**, 042006 (2001) [arXiv:gr-qc/0102012].
[19] L. A. Rawley, J. H. Taylor and M. M. Davis, Astrophys. J. **326**, 947 (1988); M. F. Ryba and J. H. Taylor, Astrophys. J. **371**, 739 (1991).
[20] P. Jaranowski and A. Krolak, Phys. Rev. D **59**, 063003 (1999) [arXiv:gr-qc/9809046].
[21] M. Zimmermann and E. Szedenits, Phys. Rev. D **20**, 351 (1979).
[22] V. M. Kaspi, J. H. Taylor and M. F. Ryba, Astrophys. J. **428**, 713 (1994).